

Table 4F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
BCA	736	91.6	1.6	4.6	0.0	0.4	1.8	0.0
BCHD	1,828	72.6	1.3	21.4	0.0	0.4	4.2	0.0
BRSI	2,565	92.6	0.6	4.2	0.1	0.3	2.2	0.0
CAP	2,359	87.7	0.8	10.2	0.0	0.1	1.2	0.0
CCHD	914	91.6	0.3	4.2	0.0	0.5	3.4	0.0
CFS	3,938	86.3	0.4	9.2	0.0	0.1	3.9	0.0
DCH	2,113	94.6	0.6	2.3	0.0	0.0	2.6	0.0
DCHD	1,017	65.0	0.4	30.9	0.1	0.6	3.0	0.0
DMH	1,084	94.7	0.6	2.6	0.0	0.8	1.3	0.0
DMHC	3,872	81.6	3.6	6.5	0.1	0.3	7.9	0.0
ECHD	2,350	2.9	32.8	62.1	0.0	0.0	2.1	0.0
ELKCHD	7,636	42.6	10.1	41.4	0.0	0.5	5.4	0.0
FHHC	304	94.7	0.0	4.9	0.0	0.0	0.3	0.0
FLCHD	1,709	65.5	13.9	9.0	0.1	3.9	7.7	0.0
FULCHD	586	85.7	0.7	11.9	0.0	0.0	1.7	0.0
GCRC	1,077	85.1	3.2	3.2	0.1	0.1	8.3	0.0
HCHF	1,447	83.8	6.4	5.1	0.1	2.8	1.7	0.0
HCMH&HS	1,001	88.5	2.0	4.6	0.2	0.0	4.7	0.0
HIS	2,235	89.1	0.7	8.7	0.0	0.0	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4F
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana

Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
HU	249	94.8	1.2	0.8	0.0	0.4	2.8	0.0
ICAP	1,286	89.7	1.1	3.8	0.1	0.4	5.0	0.0
IHC	10,949	68.0	7.2	18.4	0.1	0.7	5.6	0.0
JEFCHD	864	82.1	0.8	11.0	0.3	0.6	5.2	0.0
JNHC	1,370	93.2	0.5	2.8	0.0	0.3	3.1	0.0
JOCHD	2,558	81.2	1.8	8.1	0.0	4.1	4.8	0.0
KCHD	1,727	66.2	1.0	29.9	0.1	0.5	2.4	0.0
LCHD	610	77.7	0.7	16.6	0.0	1.1	3.9	0.0
MADCHD	3,246	68.1	12.1	12.9	0.1	0.1	6.7	0.0
MARCHD	31,659	27.1	37.7	26.9	0.1	4.1	4.0	0.0
MARSCHD	1,329	64.9	0.6	30.6	0.0	0.2	3.8	0.0
MCMH	1,450	92.8	0.1	3.4	0.0	0.1	3.5	0.0
MHS	7,045	40.3	27.6	22.7	0.1	1.2	8.2	0.0
MW	1,837	83.7	1.8	6.6	0.2	0.5	7.2	0.1
NCCAA	3,248	60.1	20.2	13.9	0.0	0.4	5.3	0.0
NHC	13,511	45.4	21.6	19.6	0.0	5.5	7.9	0.0
NHS	5,313	70.1	6.3	18.2	0.1	0.6	4.8	0.0
NWFS	17,490	39.3	34.9	22.1	0.0	0.4	3.2	0.0
ODCS	2,887	76.7	11.2	3.5	0.1	0.6	8.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4F
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana

Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
PHNA	3,191	78.3	4.5	6.4	0.0	3.4	7.3	0.0
St Mary's	2,028	89.8	0.9	4.9	0.0	0.1	4.2	0.0
STARCHD	774	91.7	0.1	6.5	0.0	0.0	1.7	0.0
STEUBCHD	1,246	87.6	0.6	8.6	0.0	1.0	2.3	0.0
SVJH	1,155	93.5	1.6	4.0	0.1	0.1	0.8	0.0
TCHD	4,738	53.3	10.4	25.8	0.2	5.0	5.3	0.0
VCHD	3,836	66.2	19.7	6.6	0.1	0.8	6.6	0.0
WCHD	746	93.8	0.4	3.9	0.0	0.1	1.7	0.0
WCMH	1,179	76.8	0.2	21.0	0.0	0.2	1.9	0.0
WICAA	3,168	78.9	7.3	4.6	0.0	0.7	8.5	0.0
WVHS	1,724	79.9	1.0	15.1	0.1	0.1	3.8	0.0
Unknown Local Ag.	5	*	*	*	*	*	*	*
Indiana	171,189	57.4	17.2	18.6	0.1	1.8	4.9	0.0
Nation (Prior Year)	8,672,965	32.5	19.1	39.7	0.9	2.5	2.7	2.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5F
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
BCA	736	36.0	19.4	44.6
BCHD	1,828	41.6	17.8	40.6
BRSI	2,565	36.6	17.2	46.2
CAP	2,359	35.6	19.4	45.0
CCHD	914	36.7	17.6	45.7
CFS	3,938	36.9	19.2	43.9
DCH	2,113	38.0	18.4	43.6
DCHD	1,017	38.0	15.9	46.1
DMH	1,084	39.6	19.4	41.1
DMHC	3,872	35.1	21.4	43.5
ECHD	2,350	35.0	19.6	45.4
ELKCHD	7,636	33.7	20.0	46.3
FHHC	304	36.8	18.8	44.4
FLCHD	1,709	37.4	21.3	41.3
FULCHD	586	32.3	21.5	46.2
GCRC	1,077	44.3	19.5	36.2
HCHF	1,447	38.5	21.1	40.4
HCMH&HS	1,001	42.0	20.3	37.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5F
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
HIS	2,235	38.9	17.9	43.2
HU	249	38.2	16.5	45.4
ICAP	1,286	34.8	19.8	45.4
IHC	10,949	36.7	18.0	45.3
JEFCHD	864	38.9	18.1	43.1
JNHC	1,370	34.1	20.8	45.1
JOCHD	2,558	42.8	17.1	40.0
KCHD	1,727	35.5	19.9	44.6
LCHD	610	33.9	17.0	49.0
MADCHD	3,246	42.9	17.9	39.2
MARCHD	31,659	34.3	23.7	42.0
MARSCHD	1,329	38.8	17.0	44.2
MCMH	1,450	44.7	14.8	40.5
MHS	7,045	35.6	21.3	43.1
MW	1,837	35.2	16.7	48.1
NCCAA	3,248	38.9	17.5	43.6
NHC	13,511	32.7	20.8	46.5
NHS	5,313	34.7	19.7	45.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5F
2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
NWFS	17,490	38.7	18.1	43.1
ODCS	2,887	41.0	18.6	40.3
PHNA	3,191	38.0	18.7	43.3
St Mary's	2,028	42.9	17.4	39.7
STARCHD	774	34.1	18.1	47.8
STEUBCHD	1,246	31.0	18.5	50.5
SVJH	1,155	33.0	18.6	48.4
TCHD	4,738	35.7	20.2	44.2
VCHD	3,836	46.0	17.3	36.7
WCHD	746	33.6	18.0	48.4
WCMH	1,179	35.5	17.6	47.0
WICAA	3,168	41.4	20.1	38.4
WVHS	1,724	32.1	21.2	46.6
Unknown Local Ag.	5	*	*	*
Indiana	171,189	36.6	19.9	43.5
Nation (Prior Year)	8,672,965	33.7	22.1	44.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
BCA	180	6.1 (4)	7.2 (31)	718	7.0 (32)	5.2 (49)	13.6 (47)	320	19.1 (44)	18.8 (47)	478	5.0 (1)
BCHD	502	8.8 (29)	5.6 (9)	1,733	6.8 (29)	1.6 (12)	10.0 (7)	733	16.2 (19)	14.2 (18)	1,132	9.4 (13)
BRSI	604	7.1 (8)	5.8 (13)	2,487	6.3 (23)	1.9 (18)	10.4 (17)	1,154	17.8 (38)	16.4 (33)	1,720	6.4 (4)
CAP	564	8.7 (25)	8.5 (43)	2,331	6.2 (18)	2.2 (22)	10.1 (9)	1,046	18.2 (39)	15.9 (27)	1,623	13.2 (28)
CCHD	213	6.1 (4)	4.7 (3)	902	7.6 (42)	1.3 (7)	12.0 (35)	407	16.7 (24)	16.7 (36)	623	13.8 (30)
CFS	947	8.2 (19)	5.4 (6)	3,903	7.2 (35)	1.9 (18)	9.9 (6)	1,717	16.5 (21)	12.3 (5)	2,639	12.3 (23)
DCH	516	9.9 (38)	8.5 (43)	2,085	5.5 (6)	1.5 (9)	10.2 (12)	913	17.3 (33)	14.3 (20)	1,472	10.7 (19)
DCHD	242	4.5 (3)	5.8 (13)	992	5.4 (5)	2.3 (27)	10.3 (16)	455	15.8 (13)	14.5 (21)	677	8.7 (12)
DMH	263	10.6 (40)	5.7 (10)	1,067	6.8 (29)	1.3 (7)	11.1 (27)	446	16.1 (17)	12.6 (7)	712	9.7 (14)
DMHC	945	7.0 (7)	8.3 (41)	3,821	6.6 (28)	2.2 (22)	10.5 (18)	1,674	17.7 (36)	13.9 (14)	2,538	18.3 (44)
ECHD	521	12.1 (46)	5.8 (13)	2,304	5.7 (11)	1.6 (12)	13.0 (43)	1,056	15.7 (11)	16.4 (33)	1,579	16.6 (39)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under-weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
ELKCHD	1,769	8.1 (18)	8.2 (40)	7,467	5.5 (6)	1.5 (9)	11.0 (25)	3,487	16.9 (29)	15.9 (27)	5,244	6.5 (5)
FHHC	74	* *	* *	301	6.0 (14)	3.3 (41)	10.0 (7)	134	13.4 (2)	13.4 (11)	216	7.4 (7)
FLCHD	431	9.5 (35)	6.3 (22)	1,644	7.4 (39)	2.1 (21)	12.7 (40)	696	20.8 (48)	17.2 (40)	1,133	13.1 (27)
FULCHD	115	4.3 (2)	6.1 (19)	575	3.3 (1)	3.5 (43)	12.3 (37)	266	14.7 (4)	16.2 (31)	429	14.2 (32)
GCRC	291	7.2 (10)	6.2 (21)	1,052	6.0 (14)	2.3 (27)	11.2 (28)	391	16.1 (17)	15.9 (27)	680	14.6 (34)
HCHF	434	9.0 (31)	5.8 (13)	1,409	7.4 (39)	4.0 (46)	9.3 (4)	578	16.8 (27)	12.3 (5)	844	21.0 (46)
HCMH&HS	275	6.9 (6)	8.7 (45)	982	9.2 (49)	1.7 (15)	13.0 (43)	373	15.3 (6)	15.8 (26)	641	12.9 (26)
HIS	560	8.6 (23)	5.7 (10)	2,183	7.6 (42)	1.1 (2)	12.7 (40)	952	16.6 (23)	17.2 (40)	1,502	10.7 (19)
HU	61	* *	* *	245	5.7 (11)	0.4 (1)	12.7 (40)	110	12.7 (1)	16.4 (33)	152	5.9 (2)
ICAP	289	7.6 (13)	7.3 (32)	1,269	6.0 (14)	3.5 (43)	10.2 (12)	579	20.2 (47)	15.2 (23)	912	11.0 (21)
IHC	2,488	8.4 (22)	5.9 (17)	10,679	6.3 (23)	2.4 (30)	10.8 (23)	4,870	15.9 (16)	14.0 (15)	7,500	16.8 (41)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under-weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
JEFCHD	204	12.3 (47)	7.4 (34)	849	7.4 (39)	1.2 (5)	13.4 (46)	353	21.0 (49)	21.2 (49)	596	8.2 (11)
JNHC	320	7.8 (15)	6.9 (28)	1,347	6.2 (18)	2.4 (30)	12.6 (38)	610	16.7 (24)	17.0 (39)	932	7.9 (10)
JOCHD	708	8.3 (20)	7.9 (39)	2,484	5.6 (9)	2.7 (36)	6.8 (1)	1,016	14.6 (3)	10.3 (2)	1,633	12.6 (24)
KCHD	415	7.7 (14)	6.3 (22)	1,698	5.8 (13)	1.2 (5)	12.6 (38)	758	18.3 (40)	18.2 (45)	1,189	7.3 (6)
LCHD	130	11.5 (43)	6.9 (28)	596	5.2 (4)	2.2 (22)	11.4 (31)	293	17.4 (34)	16.0 (30)	401	6.0 (3)
MADCHD	897	8.6 (23)	5.7 (10)	3,171	8.9 (48)	2.4 (30)	10.2 (12)	1,253	16.5 (21)	14.2 (18)	2,144	28.5 (49)
MARCHD	8,803	9.3 (32)	6.1 (19)	30,955	6.2 (18)	4.2 (47)	10.7 (21)	13,110	15.3 (6)	14.0 (15)	19,568	17.4 (42)
MARSCHD	354	4.2 (1)	7.6 (36)	1,300	7.2 (35)	1.1 (2)	13.1 (45)	581	19.4 (45)	16.2 (31)	874	7.7 (8)
MCMH	403	8.7 (25)	8.4 (42)	1,402	6.8 (29)	1.9 (18)	10.7 (21)	577	18.7 (42)	18.0 (43)	929	13.3 (29)
MHS	1,844	10.2 (39)	7.6 (36)	6,898	5.5 (6)	2.9 (39)	10.2 (12)	3,010	15.6 (10)	13.0 (9)	4,369	14.1 (31)
MW	397	7.3 (11)	6.3 (22)	1,795	6.5 (26)	1.7 (15)	8.5 (2)	875	15.7 (11)	10.6 (3)	1,237	10.2 (17)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
NCCAA	776	9.3 (32)	5.5 (8)	3,160	6.1 (17)	3.1 (40)	11.2 (28)	1,400	16.8 (27)	13.6 (12)	2,139	14.7 (35)
NHC	3,143	9.7 (37)	4.8 (4)	13,299	6.4 (25)	2.4 (30)	11.7 (34)	6,217	16.3 (20)	14.1 (17)	9,440	21.8 (47)
NHS	1,233	8.7 (25)	6.6 (27)	5,239	6.2 (18)	1.6 (12)	12.0 (35)	2,368	17.7 (36)	16.9 (38)	3,644	9.9 (16)
NWFS	4,279	8.7 (25)	6.5 (26)	17,064	6.5 (26)	2.3 (27)	10.1 (9)	7,449	14.9 (5)	11.8 (4)	11,665	14.9 (36)
ODCS	816	10.7 (41)	4.3 (2)	2,828	7.1 (33)	4.9 (48)	10.6 (19)	1,140	16.7 (24)	15.4 (24)	1,850	22.7 (48)
PHNA	753	8.9 (30)	9.0 (46)	3,134	7.1 (33)	1.7 (15)	11.0 (25)	1,372	19.5 (46)	15.6 (25)	2,139	12.7 (25)
St Mary's	516	11.6 (44)	6.0 (18)	1,970	8.6 (47)	2.7 (36)	10.8 (23)	794	15.5 (8)	13.7 (13)	1,335	17.7 (43)
STARCHD	163	8.0 (16)	7.4 (34)	748	5.6 (9)	1.1 (2)	14.4 (49)	361	15.8 (13)	18.8 (47)	570	15.3 (37)
STEUBCHD	250	8.0 (16)	11.2 (47)	1,220	4.6 (2)	2.2 (22)	9.3 (4)	626	17.1 (30)	12.8 (8)	900	10.6 (18)
SVJH	262	9.5 (35)	6.9 (28)	1,137	7.7 (44)	3.3 (41)	13.6 (47)	545	17.6 (35)	18.0 (43)	802	16.6 (39)
TCHD	1,142	7.4 (12)	7.3 (32)	4,664	4.8 (3)	2.4 (30)	10.6 (19)	2,076	17.2 (31)	13.2 (10)	3,193	14.2 (32)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
VCHD	1,076	10.8 (42)	5.2 (5)	3,727	8.0 (45)	2.2 (22)	10.1 (9)	1,386	18.8 (43)	14.8 (22)	2,428	7.8 (9)
WCHD	167	12.0 (45)	5.4 (6)	730	8.1 (46)	2.5 (35)	11.5 (33)	349	17.2 (31)	18.3 (46)	498	16.5 (38)
WCMH	269	7.1 (8)	7.8 (38)	1,158	6.2 (18)	2.8 (38)	8.6 (3)	550	15.8 (13)	10.2 (1)	807	19.6 (45)
WICAA	905	8.3 (20)	6.4 (25)	3,122	7.2 (35)	1.5 (9)	11.2 (28)	1,209	18.6 (41)	16.8 (37)	2,054	9.8 (15)
WVHS	446	9.4 (34)	3.4 (1)	1,692	7.3 (38)	3.6 (45)	11.4 (31)	787	15.5 (8)	17.7 (42)	1,080	11.9 (22)
Unknown Local Ag.	2	* *	* *	5	* *	* *	* *	2	* *	* *	3	* *
Indiana	42,957	8.9	6.4	167,541	6.4	2.6	10.8	73,424	16.4	14.3	112,865	14.8
Nation (Prior Year)	1,851,546	9.0	6.3	8,057,124	6.3	3.6	12.4	3,572,939	16.1	14.4	5,391,027	14.6

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
- (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
BCA	180	55.6 (46)	102	16.7 (22)	113	6.2 (39)	68	* *	102	13.7 (4)	335	74.9 (27)	717	48.7 (37)
BCHD	502	68.5 (23)	319	17.9 (17)	276	10.1 (16)	273	12.8 (26)	319	8.2 (24)	763	86.8 (3)	1,732	48.3 (35)
BRSI	605	74.2 (10)	528	13.8 (35)	395	7.1 (35)	431	16.9 (11)	528	6.8 (31)	1,233	82.6 (7)	2,450	35.1 (20)
CAP	566	67.3 (25)	385	14.8 (31)	405	5.9 (40)	323	15.2 (19)	385	7.0 (29)	1,110	69.7 (40)	2,284	47.8 (34)
CCHD	213	64.8 (30)	161	12.4 (42)	152	5.3 (44)	150	10.7 (37)	161	5.6 (40)	433	80.1 (14)	882	54.4 (47)
CFS	949	70.2 (17)	643	19.1 (14)	686	7.9 (28)	522	16.7 (12)	642	10.9 (10)	1,782	62.1 (47)	3,774	46.9 (33)
DCH	521	61.2 (39)	350	10.0 (45)	347	7.5 (34)	276	11.2 (33)	350	6.0 (36)	940	78.0 (17)	2,027	56.5 (48)
DCHD	242	77.3 (3)	226	23.5 (7)	173	13.9 (4)	177	15.8 (15)	226	9.7 (16)	472	75.0 (26)	967	37.3 (22)
DMH	263	67.3 (25)	219	10.0 (45)	193	3.6 (46)	164	12.2 (30)	219	6.4 (34)	460	81.7 (10)	1,013	44.2 (30)
DMHC	948	58.9 (42)	603	13.6 (36)	697	8.3 (26)	452	12.6 (27)	603	4.5 (42)	1,736	81.9 (8)	3,745	44.2 (30)
ECHD	525	62.9 (34)	459	15.0 (29)	378	11.1 (9)	362	4.7 (43)	459	3.1 (47)	1,091	65.2 (42)	2,237	11.4 (5)
ELKCHD	1,780	75.0 (8)	1,235	25.3 (4)	1,259	12.4 (6)	909	15.4 (17)	1,235	9.7 (16)	3,669	75.1 (25)	7,337	7.4 (1)
FHHC	74	* *	46	* *	51	* *	36	* *	46	* *	138	81.2 (11)	297	51.9 (43)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
FLCHD	434	64.1 (32)	228	18.9 (15)	292	10.3 (13)	188	12.2 (30)	228	7.5 (27)	739	76.3 (22)	1,651	44.8 (32)
FULCHD	115	73.9 (11)	141	14.9 (30)	116	7.8 (32)	89	* *	141	11.3 (8)	276	79.7 (15)	570	41.9 (26)
GCRC	295	52.2 (47)	279	9.3 (47)	161	5.6 (42)	217	8.3 (42)	279	5.0 (41)	403	73.0 (35)	1,007	49.2 (38)
HCHF	436	72.0 (15)	169	21.9 (9)	235	10.6 (11)	158	22.8 (3)	169	11.8 (6)	596	77.7 (19)	1,414	36.4 (21)
HCMH&HS	276	76.4 (7)	172	15.7 (27)	184	10.3 (13)	135	25.9 (1)	172	11.6 (7)	394	73.1 (34)	951	11.3 (4)
HIS	562	69.9 (18)	475	16.0 (26)	366	7.9 (28)	371	14.0 (24)	475	8.6 (23)	995	75.8 (24)	2,134	50.9 (41)
HU	61	* *	48	* *	34	* *	37	* *	48	* *	116	87.9 (2)	239	22.2 (10)
ICAP	290	69.0 (21)	220	12.7 (40)	232	6.5 (37)	161	14.3 (23)	220	9.1 (22)	606	60.2 (48)	1,232	52.9 (45)
IHC	2,502	61.4 (38)	2,204	17.5 (21)	1,895	9.0 (20)	1,680	12.6 (27)	2,204	8.0 (25)	5,093	74.8 (28)	10,480	21.8 (9)
JEFCHD	208	69.7 (19)	145	13.1 (38)	142	10.6 (11)	122	14.8 (21)	145	7.6 (26)	382	74.6 (29)	825	37.8 (23)
JNHC	321	68.5 (23)	216	14.4 (32)	243	7.0 (36)	145	15.2 (19)	216	3.7 (46)	636	74.5 (30)	1,336	56.6 (49)
JOCHD	712	70.5 (16)	605	16.5 (24)	430	10.2 (15)	474	11.0 (36)	605	6.0 (36)	1,046	74.2 (32)	2,437	28.4 (15)
KCHD	416	74.5 (9)	251	21.9 (9)	290	7.9 (28)	195	18.5 (7)	251	10.0 (13)	810	71.0 (37)	1,675	26.1 (13)

(1) Reporting period is January 1 through December 31.
 (2) TV Viewing and Smoking in Household data are not reported by all contributors.
 (3) Analyses based on one record per child.
 (4) Excludes records with unknown data or errors.
 (5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
 (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
 (8) Children aged 2 years and older included in the analyses.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
LCHD	130	73.1 (13)	123	17.9 (17)	93	* *	96	* *	123	9.8 (15)	305	83.0 (5)	583	42.5 (27)
MADCHD	905	58.3 (43)	595	17.6 (20)	545	9.9 (18)	413	18.9 (6)	595	10.6 (11)	1,315	64.0 (43)	3,066	44.1 (29)
MARCHD	8,867	62.9 (34)	3,666	26.9 (2)	5,535	11.7 (8)	2,967	11.9 (32)	3,666	6.5 (33)	9,779	63.6 (44)	28,087	24.4 (12)
MARSCHD	355	76.9 (5)	262	21.0 (12)	197	9.1 (19)	216	16.7 (12)	262	12.6 (5)	612	73.4 (33)	1,280	9.1 (2)
MCMH	405	72.6 (14)	332	16.3 (25)	203	5.9 (40)	266	17.3 (10)	332	9.6 (18)	607	62.9 (45)	1,340	50.8 (40)
MHS	1,850	77.0 (4)	910	26.9 (2)	1,195	13.4 (5)	769	15.3 (18)	910	9.3 (21)	3,128	77.7 (19)	6,860	16.8 (8)
MW	398	64.6 (31)	347	13.5 (37)	283	6.4 (38)	250	14.4 (22)	347	6.9 (30)	914	71.7 (36)	1,751	31.8 (17)
NCCAA	777	64.1 (32)	719	16.6 (23)	487	11.1 (9)	539	9.1 (39)	719	4.2 (45)	1,442	56.0 (49)	3,076	34.0 (19)
NHC	3,151	66.6 (27)	1,993	19.3 (13)	2,409	9.0 (20)	1,452	16.0 (14)	1,993	9.5 (19)	6,200	70.5 (38)	13,155	33.8 (18)
NHS	1,241	60.8 (40)	819	18.4 (16)	832	8.8 (23)	605	17.5 (9)	819	10.1 (12)	2,515	77.1 (21)	5,207	42.7 (28)
NWFS	4,294	68.6 (22)	3,809	14.4 (32)	2,954	8.7 (24)	2,874	9.5 (38)	3,809	5.8 (39)	7,794	62.9 (45)	16,640	26.1 (13)
ODCS	817	59.0 (41)	452	15.3 (28)	426	8.9 (22)	373	12.3 (29)	452	4.4 (44)	1,204	66.9 (41)	2,753	16.3 (7)
PHNA	754	76.8 (6)	656	25.2 (5)	543	15.1 (2)	508	21.3 (4)	656	16.5 (2)	1,425	89.8 (1)	3,073	23.1 (11)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
St Mary's	518	58.3 (43)	471	10.2 (44)	352	4.5 (45)	450	9.1 (39)	471	4.5 (42)	829	78.0 (17)	1,891	52.7 (44)
STARCHD	163	69.3 (20)	141	12.8 (39)	127	8.7 (24)	107	11.2 (33)	141	6.4 (34)	385	80.8 (13)	739	51.8 (42)
STEUBCHD	253	82.6 (1)	181	22.1 (8)	203	11.8 (7)	132	19.7 (5)	181	11.0 (9)	655	78.5 (16)	1,213	48.4 (36)
SVJH	262	66.4 (28)	168	14.3 (34)	164	7.9 (28)	126	18.3 (8)	168	9.5 (19)	576	76.0 (23)	1,117	50.1 (39)
TCHD	1,145	73.4 (12)	715	24.2 (6)	829	15.2 (1)	542	23.1 (2)	715	16.1 (3)	2,114	74.3 (31)	4,522	28.5 (16)
VCHD	1,082	62.6 (36)	969	10.6 (43)	690	7.7 (33)	733	11.1 (35)	969	6.7 (32)	1,452	81.9 (8)	3,601	10.1 (3)
WCHD	167	79.6 (2)	111	27.0 (1)	127	15.0 (3)	82	* *	111	18.9 (1)	373	82.8 (6)	724	41.7 (25)
WCMH	272	65.4 (29)	240	21.3 (11)	188	10.1 (16)	203	15.8 (15)	240	10.0 (13)	572	83.4 (4)	1,117	15.9 (6)
WICAA	916	61.6 (37)	538	12.6 (41)	559	5.4 (43)	395	13.7 (25)	538	7.2 (28)	1,256	70.5 (38)	3,060	53.3 (46)
WVHS	449	57.9 (45)	203	17.7 (19)	278	8.3 (26)	137	8.8 (41)	203	5.9 (38)	832	81.0 (12)	1,707	39.3 (24)
Unknown Local Ag.	2	* *	0	* *	1	* *	1	* *	0	* *	2	* *	5	* *
Indiana	43,169	66.4	28,849	18.4	28,965	9.8	22,351	13.7	28,848	7.9	72,540	71.6	161,980	30.5

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household						
Local Agency	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% (Rank)						
Nation (Prior Year)	1,689,615	63.2	908,604	25.1	1,069,942	16.9	363,039	10.7	466,279	6.1	1,200,700	79.2	4,203,554	9.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
BCA	562	8.5 (24)	5.9 (17)	2,214	7.1 (39)	4.1 (47)	11.5 (28)	951	15.1 (5)	15.2 (30)	1,503	10.4 (17)
BCHD	1,540	8.4 (22)	6.0 (19)	5,541	5.9 (13)	1.8 (15)	10.4 (12)	2,340	16.5 (18)	14.9 (25)	3,830	9.6 (12)
BRSI	1,796	9.5 (36)	5.7 (15)	7,817	7.4 (42)	2.1 (24)	10.6 (14)	3,635	17.5 (29)	15.2 (30)	5,613	9.2 (10)
CAP	1,719	8.8 (28)	7.3 (34)	7,249	6.1 (16)	2.0 (21)	10.6 (14)	3,292	18.5 (41)	16.0 (33)	5,175	13.3 (30)
CCHD	652	5.4 (1)	7.1 (31)	2,977	6.8 (32)	1.2 (3)	13.3 (46)	1,426	18.8 (43)	17.1 (41)	2,097	11.9 (21)
CFS	2,905	9.0 (30)	5.0 (6)	11,890	7.0 (35)	2.2 (28)	9.9 (7)	5,099	15.7 (8)	12.9 (8)	8,415	14.9 (34)
DCH	1,581	7.8 (12)	7.6 (39)	6,663	5.2 (7)	1.7 (12)	10.6 (14)	2,924	17.3 (27)	14.8 (24)	4,785	12.0 (22)
DCHD	736	6.7 (4)	6.7 (26)	3,014	5.6 (11)	2.2 (28)	10.2 (10)	1,310	15.8 (11)	15.1 (28)	2,181	9.1 (9)
DMH	865	10.1 (40)	5.3 (11)	3,379	7.5 (43)	1.6 (10)	11.0 (21)	1,386	16.9 (22)	13.5 (10)	2,370	8.8 (5)
DMHC	3,021	7.8 (12)	6.8 (27)	11,965	6.6 (26)	2.1 (24)	10.7 (19)	5,183	17.5 (29)	14.2 (17)	8,180	17.1 (42)
ECHD	1,714	11.0 (47)	5.7 (15)	7,351	6.3 (18)	2.1 (24)	12.7 (42)	3,359	17.1 (24)	15.1 (28)	5,245	14.2 (32)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
ELKCHD	5,542	8.1 (17)	7.7 (40)	23,897	5.3 (8)	1.4 (7)	11.6 (31)	11,097	17.6 (33)	16.1 (35)	17,289	6.6 (2)
FHHC	235	8.9 (29)	3.4 (1)	955	6.7 (30)	2.0 (21)	10.4 (12)	411	16.1 (13)	12.4 (7)	710	9.6 (12)
FLCHD	1,256	10.1 (40)	5.0 (6)	5,072	8.4 (46)	1.8 (15)	13.9 (48)	2,156	19.6 (46)	18.2 (48)	3,488	12.4 (25)
FULCHD	457	8.1 (17)	5.0 (6)	1,971	4.3 (1)	3.2 (44)	12.1 (35)	877	15.7 (8)	17.3 (44)	1,423	15.9 (38)
GCRC	854	7.7 (10)	4.2 (3)	3,300	6.6 (26)	2.0 (21)	10.6 (14)	1,324	15.3 (6)	15.2 (30)	2,243	12.8 (27)
HCHF	1,321	9.0 (30)	7.0 (30)	4,420	6.5 (23)	3.0 (42)	9.6 (4)	1,726	17.2 (26)	12.3 (6)	2,762	18.4 (45)
HCMH&HS	795	6.5 (3)	7.7 (40)	3,000	8.7 (47)	1.4 (7)	11.8 (34)	1,176	16.4 (17)	13.7 (12)	2,052	12.2 (23)
HIS	1,641	7.6 (7)	6.5 (25)	6,733	6.3 (18)	1.2 (3)	12.7 (42)	2,966	17.5 (29)	17.2 (42)	4,778	9.7 (15)
HU	173	9.2 (33)	7.5 (38)	772	5.1 (4)	0.8 (1)	10.8 (20)	333	14.7 (3)	13.5 (10)	547	4.4 (1)
ICAP	958	9.0 (30)	7.3 (34)	3,999	5.7 (12)	3.3 (46)	9.2 (2)	1,805	17.5 (29)	13.2 (9)	2,925	11.2 (20)
IHC	7,647	8.7 (27)	6.3 (23)	33,079	6.4 (22)	2.4 (31)	11.4 (27)	15,152	16.3 (15)	14.5 (21)	24,013	17.7 (44)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
JEFCHD	681	11.5 (49)	8.2 (44)	2,687	7.1 (39)	1.6 (10)	13.4 (47)	1,107	21.0 (49)	21.3 (49)	1,863	9.3 (11)
JNHC	1,021	7.8 (12)	8.7 (47)	4,288	5.9 (13)	2.8 (39)	12.5 (40)	1,909	18.6 (42)	17.2 (42)	3,013	8.8 (5)
JOCHD	2,088	7.7 (10)	7.4 (36)	7,759	5.3 (8)	2.6 (35)	7.9 (1)	3,205	15.7 (8)	11.5 (3)	5,457	11.1 (19)
KCHD	1,193	8.5 (24)	6.3 (23)	4,880	6.5 (23)	1.3 (6)	12.8 (44)	2,166	19.3 (44)	17.9 (46)	3,435	8.4 (4)
LCHD	392	9.7 (38)	5.9 (17)	1,895	6.2 (17)	1.5 (9)	11.1 (23)	923	19.6 (46)	16.5 (38)	1,351	8.8 (5)
MADCHD	2,758	8.4 (22)	5.6 (13)	9,681	9.4 (48)	2.3 (30)	11.5 (28)	3,855	16.3 (15)	14.6 (22)	6,593	23.9 (49)
MARCHD	27,249	9.7 (38)	6.0 (19)	94,859	6.3 (18)	4.1 (47)	11.1 (23)	39,247	16.1 (13)	14.0 (16)	60,365	17.6 (43)
MARSCHD	1,004	6.4 (2)	8.0 (43)	4,083	6.9 (34)	1.0 (2)	14.1 (49)	1,804	20.0 (48)	16.9 (39)	2,924	7.1 (3)
MCMH	1,206	8.1 (17)	6.9 (28)	4,462	7.0 (35)	1.7 (12)	11.6 (31)	1,816	19.4 (45)	16.1 (35)	3,103	17.0 (41)
MHS	5,595	9.5 (36)	7.1 (31)	21,762	5.1 (4)	2.9 (40)	9.7 (5)	9,711	14.6 (2)	11.4 (2)	14,413	14.8 (33)
MW	1,251	8.1 (17)	7.1 (31)	5,729	6.0 (15)	1.9 (18)	9.3 (3)	2,764	14.8 (4)	11.3 (1)	4,132	9.6 (12)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
NCCAA	2,405	10.3 (43)	5.2 (10)	9,890	6.5 (23)	2.6 (35)	12.1 (35)	4,332	17.7 (34)	13.8 (13)	6,895	12.9 (29)
NHC	9,741	9.2 (33)	5.3 (11)	40,509	6.7 (30)	2.4 (31)	11.7 (33)	18,673	16.7 (20)	13.9 (14)	28,834	19.0 (47)
NHS	3,726	8.5 (24)	5.6 (13)	15,582	6.6 (26)	1.7 (12)	12.2 (37)	6,823	17.9 (36)	17.0 (40)	11,018	12.8 (27)
NWFS	13,406	9.2 (33)	6.2 (22)	54,519	6.3 (18)	2.4 (31)	10.1 (9)	23,914	14.5 (1)	11.9 (4)	38,827	15.4 (36)
ODCS	2,478	10.5 (45)	4.4 (4)	8,820	6.6 (26)	6.1 (49)	10.3 (11)	3,548	15.9 (12)	14.4 (20)	6,015	19.5 (48)
PHNA	2,312	8.0 (16)	8.2 (44)	9,771	6.8 (32)	1.8 (15)	11.3 (26)	4,337	18.1 (37)	15.0 (27)	6,894	12.3 (24)
St Mary's	1,588	10.1 (40)	6.1 (21)	6,059	9.7 (49)	2.6 (35)	12.6 (41)	2,494	15.6 (7)	14.2 (17)	4,218	15.7 (37)
STARCHD	534	7.3 (5)	9.4 (49)	2,427	5.1 (4)	1.2 (3)	13.0 (45)	1,161	18.3 (39)	16.4 (37)	1,895	15.0 (35)
STEUBCHD	751	8.1 (17)	8.8 (48)	3,847	4.4 (2)	2.9 (40)	10.6 (14)	2,007	17.3 (27)	14.6 (22)	2,859	10.2 (16)
SVJH	789	7.6 (7)	7.7 (40)	3,589	7.6 (44)	3.1 (43)	12.2 (37)	1,748	17.0 (23)	14.9 (25)	2,642	18.8 (46)
TCHD	3,584	7.4 (6)	8.5 (46)	14,864	4.8 (3)	2.5 (34)	11.0 (21)	6,652	17.1 (24)	13.9 (14)	10,503	13.7 (31)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Local Agency	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
VCHD	3,381	10.6 (46)	4.7 (5)	12,059	7.1 (39)	1.9 (18)	9.9 (7)	4,642	17.8 (35)	14.2 (17)	8,180	8.8 (5)
WCHD	565	11.0 (47)	5.1 (9)	2,439	7.9 (45)	2.7 (38)	11.5 (28)	1,115	18.4 (40)	17.3 (44)	1,755	16.3 (39)
WCMH	838	7.9 (15)	7.4 (36)	3,636	5.4 (10)	2.1 (24)	9.7 (5)	1,695	16.6 (19)	12.2 (5)	2,665	16.7 (40)
WICAA	2,682	7.6 (7)	6.9 (28)	9,508	7.0 (35)	1.9 (18)	11.2 (25)	3,750	18.1 (37)	16.0 (33)	6,511	10.9 (18)
WVHS	1,308	10.4 (44)	3.7 (2)	5,189	7.0 (35)	3.2 (44)	12.2 (37)	2,401	16.7 (20)	18.1 (47)	3,439	12.4 (25)
Unknown Local Ag.	2	* *	* *	5	* *	* *	* *	2	* *	* *	3	* *
Indiana	132,498	9.0	6.3	522,056	6.4	2.6	11.1	227,729	16.6	14.3	361,426	14.6

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
BCA	645	57.4 (43)	453	19.9 (14)	462	9.3 (18)	439	13.4 (22)	449	10.0 (7)	1,009	73.5 (32)	1,316	46.8 (34)
BCHD	1,717	67.0 (19)	1,228	16.7 (26)	1,130	7.6 (34)	1,184	12.8 (26)	1,223	8.7 (17)	2,487	85.0 (3)	3,387	47.3 (35)
BRSI	1,988	69.6 (14)	1,667	15.1 (32)	1,525	7.7 (33)	1,581	15.0 (10)	1,661	7.0 (31)	3,896	77.2 (15)	4,350	39.4 (25)
CAP	1,912	62.2 (30)	1,449	13.7 (38)	1,443	6.0 (42)	1,408	10.4 (37)	1,442	5.8 (41)	3,519	67.7 (42)	4,099	48.4 (36)
CCHD	712	61.7 (34)	571	15.1 (32)	566	6.2 (39)	555	10.3 (39)	569	7.4 (29)	1,528	75.4 (24)	1,643	52.1 (44)
CFS	3,228	67.9 (16)	2,450	17.5 (23)	2,367	8.2 (28)	2,355	15.1 (9)	2,438	9.8 (10)	5,438	67.7 (42)	6,644	44.5 (28)
DCH	1,763	57.7 (42)	1,370	11.3 (44)	1,324	6.0 (42)	1,303	10.7 (35)	1,364	6.5 (35)	3,082	80.7 (9)	3,752	57.4 (49)
DCHD	817	74.3 (4)	702	23.5 (7)	620	11.8 (7)	674	14.4 (16)	700	8.3 (20)	1,405	76.9 (16)	1,657	37.6 (23)
DMH	930	65.2 (25)	733	11.3 (44)	677	5.9 (45)	692	10.8 (34)	732	6.1 (37)	1,475	79.3 (12)	1,865	46.0 (31)
DMHC	3,329	58.3 (41)	2,425	14.9 (35)	2,469	8.0 (32)	2,300	10.3 (39)	2,410	5.5 (43)	5,528	81.4 (8)	6,811	46.4 (33)
ECHD	1,896	61.1 (35)	1,531	17.7 (20)	1,426	10.1 (14)	1,435	4.3 (49)	1,518	2.6 (49)	3,575	61.8 (48)	4,129	12.8 (4)
ELKCHD	6,012	73.6 (6)	4,587	26.3 (3)	4,597	12.4 (5)	4,289	13.3 (23)	4,545	8.5 (18)	11,918	74.6 (27)	13,142	8.7 (2)
FHHC	258	58.5 (40)	199	7.5 (49)	190	5.3 (48)	188	9.0 (46)	198	4.0 (48)	437	74.4 (29)	549	54.1 (47)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
- (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
- (8) Children aged 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
FLCHD	1,449	64.9 (26)	1,035	19.1 (17)	1,018	8.7 (21)	1,001	12.9 (25)	1,024	7.8 (27)	2,288	73.5 (32)	2,959	46.3 (32)
FULCHD	503	66.2 (22)	424	14.6 (36)	409	6.4 (36)	392	13.5 (20)	421	8.1 (23)	922	76.6 (18)	1,118	27.6 (12)
GCRC	987	51.7 (49)	786	10.4 (48)	630	5.7 (46)	745	10.3 (39)	783	6.6 (33)	1,397	73.8 (30)	1,968	48.7 (37)
HCHF	1,493	70.1 (12)	864	22.2 (9)	876	10.3 (13)	854	19.0 (6)	856	12.9 (5)	1,835	80.5 (10)	2,798	38.8 (24)
HCMH&HS	896	70.6 (10)	616	21.4 (10)	622	10.1 (14)	607	20.8 (4)	612	14.2 (4)	1,269	73.8 (30)	1,799	16.5 (5)
HIS	1,827	65.6 (23)	1,439	17.3 (24)	1,290	8.6 (24)	1,383	14.5 (14)	1,430	8.3 (20)	3,166	74.7 (26)	3,801	51.9 (43)
HU	190	53.7 (47)	163	11.7 (43)	149	6.7 (35)	160	10.6 (36)	163	6.7 (32)	358	90.5 (1)	417	33.6 (18)
ICAP	1,031	66.9 (20)	823	16.3 (29)	817	8.2 (28)	771	14.5 (14)	819	9.9 (8)	1,923	66.6 (45)	2,174	51.5 (42)
IHC	8,541	61.8 (32)	7,109	16.7 (26)	6,602	8.6 (24)	6,692	12.4 (28)	7,077	7.9 (26)	16,229	72.3 (38)	18,342	25.3 (11)
JEFCHD	760	66.3 (21)	540	16.7 (26)	505	10.9 (11)	536	14.9 (11)	535	9.5 (12)	1,202	76.5 (19)	1,543	28.6 (15)
JNHC	1,123	64.3 (27)	849	13.8 (37)	866	8.1 (31)	799	12.8 (26)	844	4.9 (47)	2,013	75.8 (22)	2,427	53.9 (46)
JOCHD	2,276	69.7 (13)	1,826	19.2 (16)	1,539	10.0 (17)	1,723	12.4 (28)	1,816	6.6 (33)	3,362	76.1 (20)	4,581	31.3 (16)
KCHD	1,340	70.3 (11)	1,017	23.5 (7)	1,017	10.6 (12)	963	15.5 (8)	1,010	8.1 (23)	2,316	72.5 (36)	2,753	33.3 (17)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
- (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
- (8) Children aged 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
LCHD	449	72.4 (8)	378	21.4 (10)	331	14.5 (3)	356	19.7 (5)	376	11.7 (6)	973	84.8 (4)	1,077	42.5 (26)
MADCHD	2,982	56.4 (44)	2,066	17.6 (22)	1,877	10.1 (14)	1,945	13.6 (19)	2,054	9.4 (14)	4,091	68.8 (41)	5,802	45.4 (30)
MARCHD	29,389	62.0 (31)	18,365	23.6 (6)	19,894	11.7 (8)	17,653	9.6 (44)	18,154	6.0 (38)	32,140	66.7 (44)	53,829	24.1 (10)
MARSCHD	1,108	71.6 (9)	884	18.3 (18)	804	9.2 (19)	850	14.4 (16)	880	9.8 (10)	1,930	75.0 (25)	2,356	8.3 (1)
MCMH	1,332	67.6 (17)	997	15.0 (34)	835	6.3 (38)	946	14.2 (18)	994	8.4 (19)	1,935	66.0 (46)	2,652	56.0 (48)
MHS	6,136	73.1 (7)	4,081	26.5 (2)	4,313	13.7 (4)	3,924	14.8 (12)	4,043	9.2 (15)	10,341	76.1 (20)	12,650	16.7 (6)
MW	1,331	61.8 (32)	1,152	13.5 (39)	1,104	4.8 (49)	1,054	12.2 (30)	1,150	7.7 (28)	2,930	71.9 (39)	3,063	35.7 (22)
NCCAA	2,705	60.0 (37)	2,170	19.5 (15)	1,854	12.2 (6)	2,012	8.9 (47)	2,150	5.5 (43)	4,308	51.6 (49)	5,609	34.6 (20)
NHC	10,643	63.3 (29)	7,897	18.1 (19)	8,180	8.7 (21)	7,474	12.2 (30)	7,837	8.2 (22)	19,123	72.9 (34)	22,420	33.6 (18)
NHS	4,212	64.0 (28)	3,209	17.7 (20)	3,124	9.1 (20)	3,030	13.5 (20)	3,188	8.9 (16)	7,318	75.5 (23)	8,729	42.7 (27)
NWFS	14,834	67.4 (18)	11,892	15.5 (31)	10,674	8.2 (28)	11,197	9.4 (45)	11,848	5.9 (39)	25,538	64.3 (47)	31,161	27.7 (13)
ODCS	2,601	60.0 (37)	1,739	16.2 (30)	1,661	8.7 (21)	1,719	11.5 (33)	1,729	5.6 (42)	3,813	72.8 (35)	5,001	23.7 (9)
PHNA	2,589	73.8 (5)	2,100	24.8 (5)	1,937	15.5 (2)	1,993	21.4 (2)	2,077	14.8 (3)	4,577	88.3 (2)	5,503	21.7 (8)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
- (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
- (8) Children aged 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7F
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Local Agency	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
St Mary's	1,771	53.9 (46)	1,406	11.9 (42)	1,175	6.1 (40)	1,422	10.3 (39)	1,399	5.1 (46)	2,682	76.9 (16)	3,567	52.1 (44)
STARCHD	577	65.5 (24)	488	16.8 (25)	481	8.3 (26)	451	14.6 (13)	483	9.9 (8)	1,258	77.8 (14)	1,302	51.2 (41)
STEUBCHD	835	78.8 (1)	688	20.1 (12)	701	8.3 (26)	635	16.7 (7)	684	9.5 (12)	2,131	79.5 (11)	2,085	51.0 (40)
SVJH	848	61.0 (36)	666	13.2 (40)	655	6.0 (42)	602	11.6 (32)	665	8.0 (25)	1,877	78.3 (13)	1,923	50.0 (38)
TCHD	3,951	74.5 (3)	2,970	27.0 (1)	2,986	16.3 (1)	2,764	21.2 (3)	2,942	16.3 (2)	7,014	74.5 (28)	8,433	28.5 (14)
VCHD	3,777	55.1 (45)	2,907	10.9 (46)	2,474	6.1 (40)	2,730	10.3 (39)	2,899	6.4 (36)	4,971	72.4 (37)	7,245	11.1 (3)
WCHD	621	75.4 (2)	468	25.9 (4)	473	11.6 (9)	456	28.3 (1)	466	17.2 (1)	1,198	81.7 (6)	1,401	44.5 (28)
WCMH	915	68.1 (15)	771	20.0 (13)	697	11.0 (10)	730	13.2 (24)	767	7.3 (30)	1,812	84.2 (5)	1,932	17.5 (7)
WICAA	2,926	60.0 (37)	2,014	10.9 (46)	1,883	5.4 (47)	1,930	10.4 (37)	2,011	5.9 (39)	3,981	69.4 (40)	5,629	50.9 (39)
WVHS	1,444	53.3 (48)	931	12.4 (41)	1,027	6.4 (36)	879	7.5 (48)	929	5.3 (45)	2,568	81.6 (7)	3,062	35.3 (21)
Unknown Local Ag.	5	* *	3	* *	2	* *	4	* *	3	* *	2	* *	7	* *
Indiana	145,604	64.6	107,098	18.8	104,278	9.7	101,785	12.2	106,367	7.6	232,088	72.0	296,462	31.1

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.
- (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
- (8) Children aged 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

